

The effect of node-degree correlation on synchronization of identical pulse-coupled oscillators

M. Drew LaMar Gregory D. Smith

April 12, 2010

Abstract

We explore the effect of correlations between the in- and out-degrees of random directed networks on the synchronization of identical pulse-coupled oscillators. Numerical experiments demonstrate that the proportion of initial conditions resulting in a globally synchronous state (prior to a large but finite time) is an increasing function of node-degree correlation. For those networks observed to globally synchronize, both the mean and standard deviation of time to synchronization are decreasing functions of node-degree correlation. Pulse-coupled oscillator networks with negatively correlated node degree often exhibit multiple coherent attracting states, with trajectories performing fast transitions between them. These effects of node-degree correlation on dynamics of pulse-coupled oscillators are consistent with aspects of network topology (e.g., the effect of node-degree correlation on the eigenvalues of the Laplacian matrix) that have been shown to affect synchronization in other contexts.

1 Introduction

Synchronization is ubiquitous in the biological sciences. In phenomena as diverse as the synchronization of cortical neurons leading to gamma oscillations [4] and gene regulatory networks in populations of quorum sensing bacteria [15], both how the oscillators are coupled—i.e., the network structure—and the dynamics of the oscillators themselves, determine whether or not global synchronization occurs. One of the most extensively studied model systems for synchronization are the phase equations originally proposed by Kuramoto [22, 33, 1]. Interactions between Kuramoto oscillator phases occur continuously and the resulting dynamics correspond to the long-term behavior of weakly-coupled identical limit cycle oscillators.

Analysis of Kuramoto oscillators has addressed many questions regarding network structure and synchronization. Results established in the infinite- N case with all-to-all coupling have elucidated the critical coupling value for the onset to partial synchronization and the stability of the partially coherent and

fully coherent states [1]. Restrepo et al [27, 30] have considered the finite- N case with more complex network topologies, including directed networks, and established an approximate location for the critical coupling strength and local properties of the bifurcating branch. Jadbabaie et al. [19] have shown the existence of another critical coupling strength beyond which a global synchronous state exists, with its stability given by the eigenvalues of the Laplacian matrix.

Another well-studied model system for synchronization is pulse-coupled oscillators with interactions mediated by instantaneous events with finite magnitude [23, 16, 34]. Although not as realistic as integrate-and-fire or Hodgkin-Huxley-style network models, pulse-coupled oscillators are regarded as a reasonable starting point for the study of neural synchronization. While phase-response curves of realistic neuronal models have been well-studied, and synchronization of pulse-coupled oscillators have been explored in a neural context [16, 21], comparatively few studies have considered how details of network structure may influence the properties of synchronization of pulse-coupled oscillators [16, 18, 34].

This paper is focused on how a particular aspect of network structure known as “node-degree correlation” (see Sec. 2.1) affects the dynamics of synchronization of identical pulse-coupled oscillators. This question is timely because dual intracellular recording and other experimental approaches can provide data on the local connectivity of neuronal networks [26, 4, 24, 13].

2 Model formulation

We consider a population of N pulse-coupled oscillators [23] with interconnectivity given by the adjacency matrix of a directed graph, i.e., an $N \times N$ matrix A such that

$$A_{ij} = \begin{cases} 1 & \text{if } i \rightarrow j \\ 0 & \text{otherwise,} \end{cases}$$

where $i \rightarrow j$ indicates that oscillator i can affect the dynamics of oscillator j . While synchronization has been most extensively studied in the context of all-to-all coupled networks ($A_{ij} = 1$) and networks with reciprocal interactions represented by undirected graphs ($A_{ij} = A_{ji}$) [3, 10, 16], simulation and analysis of synchronization on networks represented by directed graphs are not without precedent [30, 37, 34, 19]. In this context of pulse-coupled oscillators, it is convenient to assume no self-interactions ($A_{ii} = 0$).

2.1 Network topology and node-degree correlation

Let the ordered pair $d_i = (d_i^{\text{out}}, d_i^{\text{in}})$ denote the out- and in-degree of the i -th oscillator, that is, $d_i^{\text{out}} = \sum_{j=1}^N A_{ij}$ is the number of oscillators j such that $i \rightarrow j$ and, similarly, $d_i^{\text{in}} = \sum_{j=1}^N A_{ji}$ is the number of oscillators j such that $j \rightarrow i$. The *node-degree correlation* [31] of a finite network can be quantified in

several ways, perhaps most intuitively using the Pearson correlation coefficient

$$\rho = \frac{1}{N} \sum_{i=1}^N \left(\frac{d_i^{\text{out}} - \mu^{\text{out}}}{\sigma^{\text{out}}} \right) \left(\frac{d_i^{\text{in}} - \mu^{\text{in}}}{\sigma^{\text{in}}} \right) \quad (1)$$

where μ^{out} and σ^{out} are the mean and standard deviation of the out-degree over the population of nodes (similarly for μ^{in} and σ^{in}). For example, Fig. 1(a) and (b) show two networks with $N = 6$ nodes and degree sequences given by

$$\begin{aligned} \text{(a)} \quad & \{(3, 3), (3, 3), (3, 3), (1, 1), (1, 1), (1, 1)\} \\ \text{(b)} \quad & \{(3, 1), (3, 1), (3, 1), (1, 3), (1, 3), (1, 3)\} \end{aligned} \quad (2)$$

While these two networks are similar in many ways (e.g., both have 12 edges), network (a) has a positive node-degree correlation because the Pearson correlation coefficient (Eq. 1) of its degree sequence evaluates to $\rho = 1$, while network (b) has a negative node-degree correlation ($\rho = -1$).




Figure 1: Example networks with positively (a) and negatively (b) correlated node degree. Degree sequences as in Eq. 2.

Note that the *node-degree correlation* of a directed network is distinct from the concept of *assortativity*, which in the context of undirected graphs is essentially the Pearson correlation coefficient of the degrees of pairs of connected nodes (sometimes referred to as *edge-degree correlation* [31]). Although measures of assortativity can be defined for directed networks using the in- and out-degree of nodes connected by a directed edge [38], the random networks that are the focus of this study have a wide range of node-degree correlations with negligible edge-degree correlations.

2.2 Pulse-coupled oscillators

Following [18, 23] and assuming identical intrinsic frequencies, the population of N pulse-coupled oscillators satisfies the phase equations

$$\frac{d\phi_i}{dt} = 1, \quad \text{for } i = 1, \dots, N \quad (3)$$

where $\phi_i \in [0, 1]$. Without loss of generality units of time are chosen so that the natural period of each oscillator is unity. The i -th oscillator ‘fires’ when its phase ϕ_i attains the threshold $\phi_{th} = 1$. When this occurs, the phase of the i -th oscillator is reset ($\phi_i \mapsto 0$) and the phases of oscillators along outgoing connections (j such that $i \rightarrow j$) are advanced according to

$$\phi_j \mapsto \phi_j + \begin{cases} \Delta(\phi_j) & \text{for } A_{ij} = 1 \\ 0 & \text{otherwise} \end{cases} \quad (4)$$

where $j \neq i$. We will consider three phase response curves (PRCs) $\Delta(\phi)$ – periodic on the interval $[0, 1]$ – given by

$$\begin{aligned} (a) \quad & \Delta(\phi) = \epsilon\phi, \\ (b) \quad & \Delta(\phi) = \epsilon\alpha_\theta[\sin(2\pi\phi + \theta) - \sin\theta], \\ (c) \quad & \Delta(\phi) = \epsilon\alpha_\lambda\phi(1 - \phi)e^{\lambda\phi}, \end{aligned} \quad (5)$$

where $0 \leq \theta < 2\pi$, $\lambda \geq 0$, and α_θ and α_λ are normalization constants chosen so that $\int_0^1 \Delta(\phi) d\phi = \epsilon \ll 1$. Unless otherwise noted, ϵ is given by

$$\epsilon = \frac{k}{\mu^{\text{in}}} \quad (6)$$

where $k = 0.1$ and $\mu^{\text{in}} = \frac{1}{N} \sum_{i=1}^N d_i^{\text{in}}$ is the mean in-degree. This scaling is natural for comparison across networks with different topologies, because the phase advance of the i -th oscillator with a characteristic number of incoming neighbors (j such that $j \rightarrow i$) does not depend on the network’s degree distribution when these μ^{in} neighbors fire simultaneously.




Figure 2: The θ - and λ -family of phase response curves (PRCs) (see Eq. 5). Solid and dashed lines denote curves leading to a stable and unstable synchronous state, respectively, in the case of all-to-all coupling. The significance of the labelled curves and arrows is discussed in Sec. 3.3.

The phase response curves in Eq. 5 have been chosen for different reasons. PRC 5(a) is the simplest type of excitatory interaction between oscillators that leads to a globally synchronous state for the case of all-to-all connectivity ¹.

¹Though not as physiologically relevant as the θ - and λ -family of PRCs, the linear PRC

Numerical experiments with PRC 5(a) show the synchronous state is stable for network topologies that are the focus of this paper (not shown). PRCs 5(b) (θ -family) and 5(c) (λ -family) are more relevant physiologically and satisfy

$$\Delta(0) = \lim_{\phi \rightarrow 1^-} \Delta(\phi) = 0$$

as discussed in [16] (see Fig. 2). The θ -family is similar to the PRCs for the quadratic integrate-and-fire neuron model and radial isochron clock, the latter of which includes regions of phase advance and delay. The λ -family was used in [16] to fit experimentally observed cortical neuron PRCs that are strictly phase advancing [32].

Note that the governing equations for the population of pulse-coupled oscillators used here (Eqs. 3–6) are a special case of the more general form

$$\frac{d\phi_i}{dt} = f(\phi_i) + \sum_{j=1}^N A_{ji} \Gamma(\phi_i, \phi_j) \quad (7)$$

that can be equated by choosing $f(\phi_i) = 1$, $\Gamma(\phi_i, \phi_j) = h(\phi_j)\Delta(\phi_i)$, and $h(\phi_j) = \delta(t - t_j)$, where t_j denotes the firing time of oscillator j and δ is the Dirac delta function. A finite network of all-to-all coupled Kuramoto oscillators can also be written in this form using $A_{ij} = 1$, $\Gamma(\phi_i, \phi_j) = \epsilon \sin(\phi_j - \phi_i)$, and $\epsilon = k/N$.

2.3 Network construction and sampling

In order to explore the effect of correlations between the in- and out-degrees of random directed networks on synchronization, we perform numerical experiments on strongly-connected networks whose degree sequences— $\{d_i\}_{i=1}^N$ where $d_i = (d_i^{\text{out}}, d_i^{\text{in}})$ —are drawn from a *Gaussian copula* [25], that is, a bivariate probability distribution with specified correlation ρ and discrete uniform marginals on the interval $[1, 40]$. Panels (a) and (b) of Fig. 3 show the distributions used when the node-degree correlation is $\rho = 0.7$ and -0.7 , respectively.

Random networks with specified node-degree correlation were constructed as follows. First, a random degree sequence is drawn from the appropriate Gaussian copula. Next, in- and out-degree pairs of this sequence are replaced at random with new draws from the copula until the sum of the out-degrees equals the sum of the in-degrees. If the resulting degree sequence does not have a network realization [14] a new degree sequence is drawn; otherwise, a network with this degree sequence is constructed using the algorithm of Kleitman and Wang [20]. Finally, to remove any undesired network structure, we perform a

is a simple functional form that is parameterized solely by the coupling strength ϵ . A linear PRC can be motivated using the Mirrollo-Strogatz formalism with position-phase relation $f(\phi) = \log_{a+1}(a\phi + 1)$ and the corresponding PRC given by $\Delta(\phi) = [(a+1)^{\hat{\epsilon}} - 1](\phi + \frac{1}{a})$, where $\hat{\epsilon}$ is a small positional displacement [5]. If we choose $\hat{\epsilon} = \log_{a+1}(1 + \epsilon)$, then we have the affine linear PRC $\Delta(\phi) = \epsilon(\phi + \frac{1}{a})$, which is approximated by Eq. 5(a) when a is large. In addition, PRCs for the leaky integrate-and-fire neuron model are approximately affine linear when the ratio of the oscillator period and membrane time constant is small [11].




Figure 3: Example bivariate probability distributions for in- and out-degree (d_i^{in} and d_i^{out}) used to generate random networks with specified node-degree correlation. Black and white denote high and low probability, respectively, for Gaussian copulas with uniform marginal distributions and correlation $\rho = 0.7$ (a) and -0.7 (b). Crosses in panel (a) correspond to sampled degree sequence used in Fig. 4.

Monte Carlo edge-swapping algorithm [9] and check that the resulting network is connected. This procedure well-approximates an independent random draw from the set of networks with the randomly chosen degree sequence. Simpler algorithms for constructing random graphs with specified node-degree correlation [7] would be appropriate for networks larger than those studied here.

3 Results

We will first consider results for the linear PRC in Eq. 5(a) and verify that similar qualitative results are present over a range of parameter values for the θ - and λ -family of PRCs in Eqs. 5(b) and (c), respectively. We use the standard global measure of coherence defined as

$$r(t) = \frac{1}{N} \left| \sum_{j=1}^N e^{2\pi i \phi_j(t)} \right|$$

where $r(t) \in [0, 1]$. The measure of global synchronization is small ($r \approx 0$) when the oscillators' phases are broadly distributed, but evaluates to unity ($r = 1$) when the oscillators are globally synchronized. In all simulations that follow, the initial oscillator phases are uniformly distributed on the interval $[0, 1]$ and network sizes are fixed at $N = 200$ oscillators. Qualitatively similar results are obtained for larger networks (not shown).

Figure 4 shows an example simulation of the dynamics of synchronization in a network of $N = 200$ pulse-coupled oscillators in which the correlation between in- and out-degree over the population of nodes is $\rho = 0.7$. The degree sequence of the network is illustrated by the crosses superimposed upon the Gaussian




Figure 4: (a) The stroboscopic phase ϕ of 200 pulse-coupled oscillators organized in a network with node-degree correlation of $\rho = 0.7$ (see crosses in Fig. 3(a)). Global synchronization occurs at $t = 335$ times the oscillators' natural period. (b) The corresponding coherence measure $r(t)$.

copula of Fig. 3(a). Beginning with initial oscillator phases uniformly distributed on the interval $[0, 1]$ so that $r(0) \approx 0$ (see above), a globally synchronous state is achieved at $t = 335$ (in units of the oscillators' natural period). The gradual agglomeration of clusters of oscillators is revealed by a stroboscopic plot of the evolving oscillator phases (Fig. 4(a)) as well as the standard coherence measure (Fig. 4(b)).

3.1 Node-degree correlation and global synchronization

To explore the effect of node-degree correlation on both the size of the basin of attraction for the globally synchronous state and the time until global synchrony, we repeated simulations similar to Fig. 4 for a large number of random networks with in- and out-degree correlations of $\rho \approx -1$, $\rho = 0$, and $\rho \approx 1$ (constructed as in Sec. 2.3). While all of the networks with positively correlated node degree ($\rho \approx 1$) were observed to synchronize, only 56% of networks with uncorrelated node degree ($\rho = 0$) and 3% of networks with negatively correlated node degree ($\rho \approx -1$) synchronized prior to the maximum simulation time of $t = 20,000$. Figure 5(a) shows a histogram of synchronization times for the networks with positively correlated node degree ($\rho \approx 1$) that is focused near the mean value of 460 (filled triangle). Figure 5(b) and (c) show similar histograms for the networks with uncorrelated ($\rho = 0$) or negatively correlated ($\rho \approx -1$) node degree. For those networks observed to synchronize prior to the maximum time of $t = 20,000$, the mean synchronization time is larger than that observed for networks with positively correlated node degree (Fig. 5(a)). While the mode of these distributions does not depend strongly on node-degree




Figure 5: Histograms of the synchronization time for random networks of pulse coupled oscillators with specified node-degree correlation (a) $\rho \approx 1$, (b) $\rho = 0$, and (c) $\rho \approx -1$. Filled triangles show mean synchronization time of 1,000 networks that achieved global synchrony prior to $t = 20,000$. Rightmost bar indicates percentage of networks that did not achieve global synchrony.

correlation, the heavy tails of the corresponding distributions lead to increased mean synchronization times and increased variances.




Figure 6: Mean synchronization time with 95% confidence intervals for random networks of pulse-coupled oscillators plotted as a function of node-degree correlation. Solid and dashed lines show results for initial conditions that synchronized prior to $t = 20,000$ and $40,000$, respectively.

Figure 6 shows the mean synchronization time with 95% confidence intervals for simulations as in Fig. 5 with node-degree correlations in the range $-1 < \rho < 1$. The solid and dashed lines correspond to networks observed to synchronize prior to maximum simulation times of $t = 20,000$ and $40,000$, respectively. These calculations demonstrate a general tendency for networks with positively correlated node degree to synchronize faster than networks with negatively correlated node degree. For networks with negative node-degree correlation, increasing the maximum simulation time results in larger mean synchronization times for those networks that synchronize.




Figure 7: Proportion of initial conditions leading to global synchrony prior to $t = 20,000$ (solid line and 95% confidence intervals) and $40,000$ (dashed line) as a function of node-degree correlation of random networks of pulse-coupled oscillators.

Figure 7 shows the proportion of initial conditions that lead to a globally synchronous state as an increasing function of node-degree correlation ($-1 < \rho < 1$). Furthermore, for any fixed ρ this proportion does not significantly increase when the maximum simulation time is increased from $t = 20,000$ (solid line) to $40,000$ (dashed line). Given the heavy-tailed distributions for the synchronization time of networks with negatively correlated node degree (Fig. 5(c)), it is possible that the proportion of initial conditions that lead to a globally synchronous state increases sharply for larger values of t that are not shown here. Nevertheless, the agreement of these two curves suggests that the basin of attraction for the global synchronous state is significantly smaller for networks with negatively correlated node degree.

3.2 Node-degree correlation and coherence

Figure 8 shows the synchronization measures r for networks that have not synchronized prior to the maximum simulation time of $t = 20,000$ versus node-




Figure 8: Scatter plot of synchronization measure $r(20,000)$ as a function of node-degree correlation for networks that did not synchronize prior to $t = 20,000$. The open circles and squares denote the mean and median, respectively. Inset shows distribution of synchronization measures among networks with $\rho = -0.7$.

degree correlation. The mean (open circles) and median (open squares) coherence are shown as a function of ρ . As the node-degree correlation decreases, the global synchrony measure is more broadly distributed with mean and median values of r decreasing. The fraction of networks yielding low coherence values increases dramatically as the node-degree correlation is decreased beyond $\rho \approx -0.5$. A bimodal distribution of coherence values is observed for networks with node-degree correlation in the range $-1 < \rho < -0.5$. For networks with node-degree correlation of $\rho = -0.7$, for example, the distribution of coherence values is bimodal, nearly symmetric, and has peaks at $r \approx 0.1$ and 0.9 (inset). Figure 9 shows trajectories from 3 different networks that contributed to the inset histogram in Fig. 8. Panels (a) and (b) show trajectories with a low and high coherence value, respectively, corresponding to the two modes in the histogram. Panel (c) is included to show that more interesting dynamics may occur, with the existence of two attracting coherent states (plateaus at $r \approx 0.6$ and 0.7) with multiple fast transitions between them.

Figure 10 explores the relationship between coherence and coupling strength (k in Eq. 6) in networks of pulse-coupled oscillators. Using a representative network with node-degree correlation of $\rho \approx 1$, Fig. 10(a) shows synchronization measure r (averaged over 100 trials with different initial conditions) as a function of coupling strength k at $t = 10,000$ (circles and solid line), $40,000$ (squares and dashed line) and $160,000$ (diamonds and dotted line); filled symbols indicate that all 100 initial conditions achieved global synchrony (i.e., $r = 1$). In the network with positively correlated node degree, global synchrony is obtained




Figure 9: Synchronization measure $r(t)$ for three $\rho = -0.7$ networks that did not synchronize prior to $t = 20,000$ and thus contributed to the histogram in Fig. 8 (inset).




Figure 10: The synchronization measure r , averaged over 100 trials with different initial conditions, as a function of coupling strength k (see Eq. 6) for representative networks with node-degree correlation of (a) $\rho \approx 1$, (b) $\rho = 0$ and (c) $\rho \approx -1$. Snapshots are shown at three times: $t = 10,000$ (circles and solid line), $40,000$ (squares and dashed line), and $160,000$ (diamonds and dotted line). Filled symbols indicate global synchrony ($r = 1$) over all initial conditions.

prior to $t = 10,000$ regardless of initial conditions so long as the coupling is sufficiently strong ($k > 3 \times 10^{-3}$).

Figure 10(b) and (c) show similar results obtained using representative net-

works with node-degree correlation $\rho = 0$ and $\rho \approx -1$, respectively. A comparison of solid, dashed and dotted lines shows that when oscillators are weakly coupled the networks exhibit long transients with low average coherence, followed by a transition to larger average coherence, the value of which is insensitive to the coupling strength k . In particular, the plateaus in Fig. 10(c) suggest the co-existence of high coherence asymptotically stable attracting states and low coherence states that can trap trajectories for a long period of time before releasing them. Perhaps most importantly, Fig. 10(c) suggests that coupling strengths required for global synchronization are relatively large for random networks with negatively correlated node degree.




Figure 11: Scatter plots of the synchronization measure r across trials with different initial conditions for networks with negatively correlated node degree ($\rho \approx -1$) and a range of coupling strengths k . Open squares correspond to average values shown in Fig. 10(c). The arrows at $k = 0.1$ correspond to Fig. 12. Snapshots are shown at three times: $t = 10,000$ (a), $40,000$ (b) and $160,000$ (c).

Figure 11 shows scatter plots of the synchronization measure r for networks with negatively correlated node degree ($\rho \approx -1$); the average values (open squares) in Fig. 11(a), (b) and (c) correspond to the circles, squares and diamonds, respectively, in Fig. 10(c). As the coupling strength k is varied, plateaus in the average synchronization measure often coincide with clusters of points indicating relatively low variance in the synchronization measure across simulations that are identical save for random initial phases of the oscillators. For values of the coupling strength in transitional regions between plateaus, the synchronization measures observed are more broadly distributed and, for coupling strengths in the approximate range $10^{-2} \leq k \leq 10^{-1}$, two distinct clusters are observed. Comparison of Fig. 11(a), (b), and (c) (corresponding to $t = 10,000$, $40,000$ and $160,000$, respectively) shows that these clusters are long-lived.

Figure 12 shows an example time course for a network with negative node-

degree correlation ($\rho \approx -1$) and coupling strength of $k = 0.1$ (corresponding to arrows in Fig. 11). The synchronization measure r indicates low coherence until $t \approx 20,000$, at which point the coherence abruptly increases. Qualitatively similar results were obtained for multiple random initial conditions. Among these simulations, the time of the abrupt increase in coherence varied widely, while the time-averaged coherence value during the plateaus was nearly constant.




Figure 12: Time course of synchronization measure $r(t)$ for a network with negatively correlated node degree ($\rho \approx -1$) and coupling strength $k = 0.1$ (cf. arrows in Fig. 11). Insets show distribution of oscillator phases at $t = 10,000$ and $30,000$.

3.3 Physiologically realistic PRCs

We now consider whether node-degree correlation effects the synchronization properties of pulse-coupled networks with physiologically realistic PRCs, specifically, the θ - and λ -family presented in Fig. 2.

Figure 13 (a) and (b) show the mean synchronization measure $r(t = 20,000)$ for 100 node-degree correlated networks ($\rho \approx 1$) as a function of the coupling strength k and the PRC shape parameters θ and λ , respectively. Using the conditions presented in [16], the dashed curves separate regions of stability and instability of the synchronous state under the assumption of all-to-all connectivity (see legend). For the θ family of PRCs, the high coherence region is contained within the stable region ($\pi/2 \leq \theta \leq 3\pi/2$), in spite of the fact that the networks are not all-to-all coupled. This is not the case for the λ family of PRCs, for which the stable region is to the right of the dashed curve.

To illustrate the effect of node-degree correlation on synchronization, Fig. 13 (c) and (d) give the absolute difference of the mean synchronization measure $r(t = 20,000)$ calculated from networks with $\rho \approx 1$ and $\rho \approx -1$. Consistent




Figure 13: Top panels show mean synchronization measure $r(t = 20,000)$ for 100 node-degree correlated networks ($\rho \approx 1$) as a function of the coupling strength k and the PRC shape parameters (a) θ and (b) λ (see Fig. 2). Bottom panels show the absolute difference of the mean synchronization measure $r(t = 20,000)$ calculated from networks with $\rho \approx 1$ and $\rho \approx -1$. The dashed curves identify regions of stability for the synchronous state under the assumptions of all-to-all connectivity: $\pi/2 \leq \theta \leq 3\pi/2$ (exact), $\lambda > 5.25$ (approximate) [16].

with results presented above using the linear PRC, there are parameter regions for the θ - and λ -family of PRCs for which the synchronization properties of networks with anti-correlated node degree are markedly different from networks with correlated node-degree. The PRCs that maximize this difference when the coupling strength is $k = 0.1$ are shown in Fig. 2 ($\theta \approx 2\pi/3$, $\theta \approx 4\pi/3$, $\lambda = 3.5$).

To explore more fully the effect of node-degree correlation on synchronization in the context of these physiologically realistic PRCs, Fig. 14 (a) and (b) show scatter plots of the time to synchronization for networks that synchronized prior to $t = 20,000$ ($\theta \approx 4\pi/3$ and $\lambda = 3.5$, respectively). Mean values are denoted by open circles and are omitted for node-degree correlations that did not result in at least 200 sample points. Consistent with results presented above using the linear PRC, decreasing node-degree correlation is associated with longer synchronization times. Numerical evidence suggests the synchronous state is stable for all simulations using the θ -family PRC with $\theta \approx 4\pi/3$. However, for the λ -family PRC with $\lambda = 3.5$ there is an increased proportion of simulations resulting in an unstable synchronous state as the node-degree correlation decreases (not shown).

For networks that did not synchronize prior to $t = 20,000$, Fig. 14 (c) and (d) show that decreasing node-degree correlation is associated with smaller coherence for the representative θ - and λ -type PRCs. Bimodal distributions of




Figure 14: Top panels show scatter plots of the time to synchronization for networks that synchronized prior to $t = 20,000$ using (a) $\theta \approx 4\pi/3$ and (b) $\lambda = 3.5$ (see Fig. 2). Open circles denote mean values and are omitted for node-degree correlations that did not result in at least 200 sample points. Bottom panels show scatter plots of $r(t = 20,000)$ for those networks that did not synchronize prior to $t = 20,000$. The coupling strength $k = 0.1$.

the coherence measure were observed for the λ -type PRC with node-degree correlation in the range $-0.4 \leq \rho \leq -0.2$ (not shown), similar to the linear PRC (see inset of Fig. 8). While the coherence is a gradually increasing function of node-degree correlation for the λ -type PRC (similar to the linear PRC), r is not a strong function of ρ for the θ -type PRC. It is unclear what aspect of these representative PRCs is responsible for this difference.

Figure 15 shows two representative time courses of the synchronization measure $r(t)$ for networks with negatively correlated node-degree ($\rho \approx -1$) and physiologically realistic PRCs. Figure 15 (a) shows a long-lived low coherent state followed by an abrupt transition to global synchrony at $t \approx 11,000$ ($\theta = 4\pi/3$, $k = 0.275$). Figure 15 (b) shows low- and high-coherence attracting states with many abrupt transitions between them ($\lambda = 3.5$, $k = 0.275$). In both cases qualitatively similar results were obtained using the linear PRC (cf. Fig. 9(c) and 12).

4 Discussion

We investigated the dynamics of synchronization in random networks of pulse-coupled oscillators with specified node-degree correlation ($-1 < \rho < 1$). We found that networks with negatively correlated node degree are less likely to




Figure 15: Representative time courses of the synchronization measure $r(t)$ for networks with negatively correlated node-degree correlation ($\rho \approx -1$), coupling strength $k = 0.275$, and (a) $\theta = 4\pi/3$ and (b) $\lambda = 3.5$ (see Fig. 2).

achieve global synchrony (Figs. 5 and 14) and synchronize more slowly (Fig. 6 and 14) than networks with positively correlated node degree. In networks with negatively correlated node degree, the proportion of initial conditions leading to global synchrony is reduced (Fig. 7 and 14) and the variability of the coherent state is increased (Fig. 8 and 14) in comparison to networks with uncorrelated or positively correlated node degree. Networks with negatively correlated node degree often exhibited interesting dynamics, such as long-lived transients and fast transitions between multiple attracting states, phenomena that were not observed in networks with positively correlated node degree (Figs. 9(c), 11, 12, and 15).

As mentioned in Sec. 2.1, the node-degree correlation of a directed network is not to be confused with assortativity or edge-degree correlation, concepts most often applied in the context of undirected graphs. While measures of assortativity can be defined for directed networks, the random networks studied here are not significantly edge-degree correlated. While edge-degree correlation of non-linear oscillators has been studied in the context of Laplacian dynamics [10], to our knowledge this is the first study of the effect of node-degree correlation in random networks of pulse-coupled oscillators.

4.1 Initial oscillator phase distribution and synchronization time

In all simulations presented above, the initial oscillator phases were uniformly distributed on the interval $[0, 1]$ leading to small initial coherence ($r(0) \approx 0$).

When networks with negatively correlated node degree are initialized in this fashion, global synchrony is unlikely and synchronization, when it does occur, is slow (Figs. 7 and 14). Nevertheless, it is possible to choose initial conditions with small initial coherence ($r(0) \approx 0$) that lead to rapid synchronization. For example, if the ordering of initial oscillator phases ($\phi_n(0) \geq \phi_{n+1}(0)$) coincides with the lexicographical ordering of the node degrees ($d_n^{\text{out}} \geq d_{n+1}^{\text{out}}$ and $d_n^{\text{out}} = d_{n+1}^{\text{out}} \Rightarrow d_n^{\text{in}} \geq d_{n+1}^{\text{in}}$), the time to synchronization of networks with $\rho \approx -1$ is shorter than that observed for networks with $\rho \approx 1$ when using the linear and θ -family PRCs ($\theta \approx 4\pi/3$, not shown). We found no such effect for the λ -family of PRCs when $\lambda = 3.5$, presumably because the globally synchronous state is unstable (Fig. 13).

4.2 Node-degree correlation and global network topology

The Laplacian of an undirected network with adjacency matrix A is defined as $L = D_A - A$, where D_A is the diagonal matrix of row sums of A . The eigenvalues of L give information about the network topology, such as the number of connected components, the community structure, and the rate of mixing of random walkers [8]. In the case of Kuramoto oscillators coupled via an undirected network, the Laplacian provides information about the time to synchronization [19, 2]. Although such a link has yet to be shown for directed networks of pulse-coupled oscillators, both the undirected and directed Laplacian (defined below) provide information about network topology that may be relevant to synchronization of pulse-coupled systems.

The Laplacian of a strongly-connected directed network can be defined as [6, 36]

$$L' = D_\phi - \frac{D_\phi P + P^T D_\phi}{2} \quad (8)$$

where $P = D_A^{-1}A$, A is the adjacency matrix for the network, D_A is the diagonal matrix of row sums of A , D_ϕ is a diagonal matrix, and ϕ solves $\phi P = \phi$ subject to $\|\phi\|_1 = 1$. Since L' is symmetric and has zero row sum, the eigenvalues of L' are real, nonnegative, and can be ordered as $0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_N$.

The spectral gap of the Laplacian of a directed network (λ_2) has properties similar to the spectral gap of the Laplacian of an undirected network [36]. The spectral gap is known to have an effect on the rate of convergence to the globally synchronous state when identical oscillators are coupled as in the Kuramoto model or via an undirected Laplacian ($d\phi_i/dt = \omega - \sigma \sum_j L_{ij}\phi_j$) [19, 2]. Similarly, the eigenvalue ratio λ_N/λ_2 is related to the propensity of identical oscillators coupled via an undirected Laplacian to have a stable synchronous state [3]. While it is not known if the directed Laplacian (Eq. 8) has these properties in the case of pulse-coupled oscillators, it is of interest to explore the effect of node-degree correlation on both λ_2 and the ratio λ_N/λ_2 .

Figure 16 shows the mean Laplacian eigenvalue λ_2 (panel a, bottom line) and the ratio λ_N/λ_2 (panel b) are monotone increasing and decreasing functions of node-degree correlation, respectively (average over 1,000 random networks

for each ρ). Recall that for pulse-coupled networks with negatively correlated node degree (corresponding to small λ_2 and large λ_N/λ_2 in Fig. 16) we observe slow synchronization (Fig. 6) and reduced percentage of initial conditions that reached global synchronization (Fig. 7). This relationship between Laplacian eigenvalues and synchronization is consistent with prior work focusing on oscillators coupled as in the Kuramoto model or via an undirected Laplacian [19, 2, 3]. Note this correspondence is less clear in the case of the λ -family PRCs where the synchronous state may be unstable (Fig. 14(b) and (d)).




Figure 16: Laplacian eigenvalues λ_N (top line) and λ_2 (bottom line) in (a) and ratio λ_N/λ_2 in (b) as a function of node-degree correlation (ρ). Each point shows mean $\pm$ standard deviation for 1,000 random networks.




Figure 17: Three measures of network topology as a function of node-degree correlation: (a) average shortest path length (left axis, decreasing function) and average clustering coefficient (right axis, increasing function); and (b) average communicability. Each point shows mean $\pm$ standard deviation for 1,000 random networks.

Figure 17 shows the effect of node-degree correlation on three additional measures of network topology: average shortest path length, clustering coefficient, and communicability [12]. Figure 17(a) shows that the average shortest path length and clustering coefficient are decreasing and increasing functions of node-degree correlation, respectively, consistent with the known relationship between these measures and synchronization of Kuramoto oscillators on undirected scale-free networks [17]. Figure 18 sheds light on the observed relationship be-

tween node-degree correlation and these measures by comparing the adjacency matrices for networks of 200 lexicographically ordered oscillators (Sec. 4.1) with $\rho \approx 1$ and $\rho \approx -1$. In particular, the asymmetry present in networks with negatively correlated node degree gives a directional bias to network connectivity that leads to an increased average distance between oscillators and reduced clustering.




Figure 18: Adjacency matrices for networks with node-degree correlations (a) $\rho \approx 1$ and (b) $\rho \approx -1$ with lexicographically ordered oscillators (see Sec. 4.1).

4.3 Local network topology and the dynamics of synchronization

To obtain more insight into the dynamics of synchronization in positively and negatively correlated networks, we considered local network properties. For example, Fig. 19 shows how the node-degree correlation of a network influences local clustering of subnetworks composed of nodes with similar in-degree. Local clustering of a subnetwork is defined as the number of arcs divided by the total possible number of arcs. For both positive and zero correlated node-degree networks, we find the local clustering increases with in-degree. However, for negatively correlated networks, local clustering is a biphasic function of in-degree with maximum at $d^{\text{in}} \approx 20$. This topological difference between positively and negatively correlated node-degree networks influences the dynamics of synchronization (see below).

Figure 20(a) shows the coherence measure as a function of time calculated on the subnetworks defined above. In this positively correlated network ($\rho \approx 1$) that achieves synchrony, the larger in-degree nodes synchronize first, consistent with these subnetworks having larger in-degree and larger clustering as in Fig. 19 (solid line). Using the negatively correlated network ($\rho \approx -1$) that led to the jump seen in Fig. 12, Fig. 20(b) shows higher coherence occurring instead in the subnetworks with $d^{\text{in}} \approx 20$, as well as slightly higher coherence occurring in the subnetworks with higher in-degree (cf. Fig 19). Note that in both cases, the subnetworks with relatively large clustering and large in-degree have higher coherence.




Figure 19: Average local clustering of subnetworks indexed by in-degree for node-degree correlations of $\rho \approx 1$ (solid), $\rho = 0$ (dotted), and $\rho \approx -1$ (dashed). For each d_i^{in} in the range $3 \leq d_i^{\text{in}} \leq 38$, a subnetwork was defined by selecting all nodes with in-degrees in the window $[d_i^{\text{in}} - 2, d_i^{\text{in}} + 2]$. Error bars suppressed for clarity.




Figure 20: Synchronization measure $r(t)$ calculated on subnetworks from Fig. 19 illustrating (a) synchronization on a positively correlated network ($\rho \approx 1$) and (b) the dynamics occurring during the jump illustrated in Fig. 12.

4.4 Connections with previous work

The long-lived transient states observed in networks with negatively correlated node degree (e.g., the $t = 10,000$ inset of Fig. 12) were insensitive to random perturbations of oscillator phases, that is, random perturbations did not immediately lead to a transition from the low-coherence to the high-coherence state, although it does of course change the time at which the jump ultimately occurs (not shown). In prior work that did not consider node-degree correlation, long-lived transient states were observed in networks of identical pulse-coupled oscillators with and without time delay [35, 40, 39].

Similar to the results reported here, node-degree correlation has been shown to influence the synchronization of Kuramoto oscillators whose governing equa-

tion is given by Eq. 7 with $\Gamma(\phi_i, \phi_j) = \epsilon \sin(\phi_j - \phi_i)$ and $\epsilon = k/N$ [28, 27, 29]. In particular, the onset of synchronization is shifted to smaller values of the coupling strength in directed networks of Kuramoto oscillators with positively correlated node degree [30, 31]. Restrepo and coworkers have shown that node-degree correlation is proportional to the largest eigenvalue of the network adjacency matrix (so long as the largest degree is not too large) and, in the case of Kuramoto oscillators, the critical coupling strength is inversely proportional to the largest eigenvalue. Our observations of the effect of coupling strength and node-degree correlation on synchronization of pulse-coupled oscillators similarly suggest a tendency of the positively correlated networks to synchronize more easily than negatively correlated networks (Fig. 10).

Low-coherence states tend to be associated with heterogeneity in networks of coupled oscillators. For example, synchronization is accelerated in all-to-all coupled networks of Kuramoto oscillators when variability of intrinsic frequency is reduced. Similarly, in random networks of pulse-coupled oscillators, synchronization is promoted by scaling the efficacy of connections so that each oscillator receives the same input when its incoming neighbors fire [35, 40]. As expected, when this scaling was applied to our networks of pulse-coupled oscillators, mean synchronization time and percent synchronization are no longer strong functions of node-degree correlation. Nevertheless, our observations of a strong effect of node-degree correlation on the dynamics of synchronization (Figs. 6–8, 14) is not an obvious consequence of heterogeneous network structure, because in our simulations networks with positively and negatively correlated node degree are equally heterogeneous, e.g., the marginal distribution of in- and out-degree is identical in all simulations performed.

Acknowledgments

The authors would like to thank the referees and editor for useful comments and suggestions. This material is based upon work supported by the NSF under Grant No. 0443843 to Smith. LaMar was funded by Howard Hughes Medical Institute Science Education Grant 52005868 to The College of William and Mary. This work was performed using computational facilities enabled by grants from Sun Microsystems and Virginia’s Commonwealth Technology Research Fund.

References

- [1] Juan Acebrón, L Bonilla, Conrad Vicente, Felix Ritort, and Renato Spigler. The kuramoto model: A simple paradigm for synchronization phenomena. Rev. Mod. Phys., 77(1):137–49, Jan 2005.
- [2] Juan Almendral and Albert Díaz-Guilera. Dynamical and spectral properties of complex networks. New Journal of Physics, 9(6):187–187, 2007.

- [3] Mauricio Barahona and Louis M Pecora. Synchronization in small-world systems. Phys. Rev. Lett., 89(5):054101, Jul 2002.
- [4] M Bazhenov, N F Rulkov, and I Timofeev. Effect of synaptic connectivity on long-range synchronization of fast cortical oscillations. J Neurophysiol, 100(3):1562–75, Sep 2008.
- [5] Henk Broer, Konstantinos Efsthathiou, and Easwar Subramanian. Heteroclinic cycles between unstable attractors. Nonlinearity, 21(6):1385–1410, 2008.
- [6] Fan Chung. Laplacians and the cheeger inequality for directed graphs. Annals of Combinatorics, 9(1):1–19, Apr 2005.
- [7] Fan Chung, Linyuan Lu, and Van Vu. Spectra of random graphs with given expected degrees. Proc. Natl. Acad. Sci. U.S.A., 100(11):6313–8, May 2003.
- [8] Fan R. K. Chung. Spectral Graph Theory (CBMS Regional Conference Series in Mathematics, No. 92). American Mathematical Society, February 1997.
- [9] George W Cobb and Yung-Pin Chen. An application of markov chain monte carlo to community ecology. The American Mathematical Monthly, 110(4):265–288, 2003.
- [10] M di Bernardo, F Garofalo, and F Sorrentino. Effects of degree correlation on the synchronization of networks of oscillators. Int. J. Bifurcation Chaos Appl. Sci. Eng., 17(10):3499 – 3506, 2007.
- [11] Albert Díaz-Guilera, Conrad Pérez, and Alex Arenas. Mechanisms of synchronization and pattern formation in a lattice of pulse-coupled oscillators. Phys. Rev. E, 57(4):3820, Apr 1998.
- [12] Ernesto Estrada and Desmond J Higham. Network properties revealed through matrix functions. University of Strathclyde Mathematics Research Report, 17:1–22, Mar 2008.
- [13] Danny Eytan and Shimon Marom. Dynamics and effective topology underlying synchronization in networks of cortical neurons. J. Neurosci., 26(33):8465–76, Aug 2006.
- [14] Delbert Ray Fulkerson. Zero-one matrices with zero trace. Pacific J. Math., 10(3):831–836, Mar 1960.
- [15] Jordi Garcia-Ojalvo, Michael B Elowitz, and Steven H Strogatz. Modeling a synthetic multicellular clock: repressilators coupled by quorum sensing. Proc. Natl. Acad. Sci. U.S.A., 101(30):10955–60, Jul 2004.
- [16] Pranay Goel and Bard Ermentrout. Synchrony, stability, and firing patterns in pulse-coupled oscillators. Physica D: Nonlinear Phenomena, 163(3-4):191–216, Mar 2002.

- [17] Jesús Gómez-Gardeñes and Yamir Moreno. Synchronization of networks with variable local properties. International Journal of Bifurcation & Chaos in Applied Sciences & Engineering, 17(7):2501–2507, 2007.
- [18] X Guardiola, Albert Díaz-Guilera, M Llas, and Conrad Pérez. Synchronization, diversity, and topology of networks of integrate and fire oscillators. Phys. Rev. E, 62(4):5565–5570, Oct 2000.
- [19] A Jadbabaie, N Motee, and M Barahona. On the stability of the kuramoto model of coupled nonlinear oscillators. In Proceedings of the American Control Conference, volume 5, pages 4296– 4301, 2004.
- [20] D Kleitman and D Wang. Algorithms for constructing graphs and digraphs with given valences and factors. Discrete Math., 6(1):79–88, 1973.
- [21] Tae-Wook Ko and G. Bard Ermentrout. Phase-response curves of coupled oscillators. Phys. Rev. E, 79(1):016211, Jan 2009.
- [22] Yoshiki Kuramoto. Chemical oscillations, waves, and turbulence. Springer, 1984.
- [23] Renato Mirollo and Steven Strogatz. Synchronization of pulse-coupled biological oscillators. SIAM J. Appl. Math., 50(6):1645–1662, 1990.
- [24] Gábor Molnár, Szabolcs Oláh, Gergely Komlósi, Miklós Füle, János Szabadics, Csaba Varga, Pál Barzó, and Gábor Tamás. Complex events initiated by individual spikes in the human cerebral cortex. PLoS Biol, 6(9):1842–1849, Sep 2008.
- [25] Roger B. Nelsen. An Introduction to Copulas. Springer, 1998.
- [26] Anne-Marie M Oswald and Alex D Reyes. Maturation of intrinsic and synaptic properties of layer 2/3 pyramidal neurons in mouse auditory cortex. J Neurophysiol, 99(6):2998–3008, Jun 2008.
- [27] Juan G Restrepo, Edward Ott, and Brian R Hunt. Onset of synchronization in large networks of coupled oscillators. Phys. Rev. E, 71(3 Pt 2A):036151, Mar 2005.
- [28] Juan G Restrepo, Edward Ott, and Brian R Hunt. Characterizing the dynamical importance of network nodes and links. Phys. Rev. Lett., 97(9):094102, Sep 2006.
- [29] Juan G Restrepo, Edward Ott, and Brian R Hunt. Emergence of coherence in complex networks of heterogeneous dynamical systems. Phys. Rev. Lett., 96(25):254103, Jun 2006.
- [30] Juan G Restrepo, Edward Ott, and Brian R Hunt. Synchronization in large directed networks of coupled phase oscillators. Chaos, 16(1):015107, Mar 2006.

- [31] Juan G Restrepo, Edward Ott, and Brian R Hunt. Approximating the largest eigenvalue of network adjacency matrices. Phys. Rev. E, 76(5):056119, 2007.
- [32] A D Reyes and E E Fetz. Two modes of interspike interval shortening by brief transient depolarizations in cat neocortical neurons. J Neurophysiol, 69(5):1661–72, May 1993.
- [33] Steven H Strogatz. From kuramoto to crawford: exploring the onset of synchronization in populations of coupled oscillators. Physica D: Nonlinear Phenomena, 143(1-4):1–20, Sep 2000.
- [34] Marc Timme. Collective dynamics in networks of pulse-coupled oscillators. Collective dynamics in networks of pulse-coupled oscillators (PhD Thesis), Oct 2002.
- [35] Marc Timme, Fred Wolf, and Theo Geisel. Prevalence of unstable attractors in networks of pulse-coupled oscillators. Phys. Rev. Lett., 89(15):154105, Sep 2002.
- [36] Chai Wu. On rayleigh-ritz ratios of a generalized laplacian matrix of directed graphs. Linear Algebra and Its Applications, 402:207–227, Jun 2005.
- [37] Chai Wu. Synchronization in networks of nonlinear dynamical systems coupled via a directed graph. Nonlinearity, 18(3):1057–1064, 2005.
- [38] Gorka Zamora-López, Changsong Zhou, Vinko Zlatić, and Jürgen Kurths. The generation of random directed networks with prescribed 1-node and 2-node degree correlations. Journal of Physics A: Mathematical and Theoretical, 41(22):224006, 2008.
- [39] R Zillmer, Roberto Livi, Antonio Politi, and Alessandro Torcini. Desynchronization in diluted neural networks. Phys. Rev. E, 74(3 Pt 2):036203, Sep 2006.
- [40] Alexander Zumdieck, Marc Timme, Theo Geisel, and Fred Wolf. Long chaotic transients in complex networks. Phys. Rev. Lett., 93(24):244103, 2004.